Provided for non-commercial research and educational use only. Not for reproduction or distribution or commercial use.



This article was originally published in a journal published by Elsevier, and the attached copy is provided by Elsevier for the author's benefit and for the benefit of the author's institution, for non-commercial research and educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are prohibited. For exceptions, permission may be sought for such use through Elsevier's permissions site at:

http://www.elsevier.com/locate/permissionusematerial



Available online at www.sciencedirect.com



GEODERMA

Geoderma 136 (2006) 609-620

www.elsevier.com/locate/geoderma

Formation and persistence of Mollisols on zeolitic Deccan basalt of humid tropical India

T. Bhattacharyya *, D.K. Pal, S. Lal, P. Chandran, S.K. Ray

Division of Soil Resource Studies, National Bureau of Soil Survey and Land Use Planning, Amravati Road, Nagpur 440 010, Maharashtra, India

Received 8 September 2005; received in revised form 21 April 2006; accepted 25 April 2006 Available online 30 June 2006

Abstract

In contrast to generally observed non-acidic and less weathered Mollisols in temperate semi-arid and humid climate, acidic and fairly weathered Mollisols on Deccan basalt are observed in hills of central (Satpura Range, Madhya Pradesh) and Western India (Western Ghats, Maharashtra) under forest in the present tropical humid climatic conditions. The detailed morphological, chemical and mineralogical investigations of these Mollisols of central and western India indicates that due to the presence of zeolites in the Deccan basalt, the transformation of smectite to kaolinite was prevented and the retention of adequate amount of smectite and continuous supply of calcium ions from zeolites made the formation of Mollisols in forest even under tropical humid climate.

The formation and the persistence of these Mollisols in association with acidic Alfisols under sparse forest in the basaltic landscape of millions of years in central and western India demonstrates the primary importance of the quality of parent material rich in expanding clay mineral and bases in their formation. This unique example thus expands the basic knowledge on the formation of Mollisols in the humid tropical part of the world.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Mollisols; Tropical humid climate; Zeolite

1. Introduction

Mollisols represent dark coloured, base-rich (*mollic* epipedon) soils of the steppes and cover extensive subhumid to semi-arid areas of north and south America, Europe and Asia. Most of these soils have a grass vegetation after they were deforested. By and large Mollisols occur in landscapes covered by grasslands but they also occur in forested areas. In high altitudes, Mollisols have formed in late Pleistocene or Holocene deposits. Beyond the limits of glaciation, Mollisols may

* Corresponding author. Fax: +91 712 2500534. *E-mail addresses:* tapas11156@yahoo.com, tapas@nbsslup.ernet.in (T. Bhattacharyya). be in older deposits or on older surfaces dating back perhaps to mid-Pleistocene or earlier (Soil Survey Staff, 1999). Mollisols in the USA have formed mostly in Quaternary materials on gentle or moderate slopes. They occur in a wide range of landscapes ranging from flat alluvial plains to undulating plains and mountains (Fenton, 1983).

The major intention of defining Mollic epipedon (of Mollisols) was to provide a differentia to distinguish the soils that have traditionally been used to produce grain, from those that are too dry to cultivate without irrigation (Smith, 1983). The examples of the soils reported from the temperate world even in agricultural land use system may contain high organic matter with soft (*mollus*) soil structure to be termed as Mollisols due to cold climate

^{0016-7061/\$ -} see front matter 0 2006 Elsevier B.V. All rights reserved. doi:10.1016/j.geoderma.2006.04.021

preventing loss of organic matter. But in the humid tropics, the soils under agricultural lands are subjected to high atmospheric temperature and therefore can not store enough organic matter to be grouped under Mollisols. Interestingly, Mollisols have been reported under subtropical part of northern India developed in micaceous alluvium of the Indo-Gangetic Plains (Deshpande et al., 1971; Murthy et al., 1982). These soils are formed under environments similar to temperate climate of the USA and Europe. More recently Mollisols have been reported from hilly mountains of southern Peninsular and north-eastern India (Krishnan et al., 1996; Das et al., 1996; Shivaprasad et al., 1998). On the high altitude plateau surfaces of central and western India under humid tropical climate, highly weathered ferruginous soils (Ultisols/or Oxisols) are expected to form; but in reality these ferruginous soils are base-rich Alfisols. A positive role of zeolites of amygdoloidal Deccan basalt in the persistence of these non-kaolinitic and/or non-oxidic Alfisols in this climate since the early Tertiary was demonstrated by Bhattacharyya et al. (1999). In addition, instead of commonly reported neutral to alkaline Mollisols, occurrence of acidic Mollisols under forest associated with Alfisols and Vertisols in the humid tropics of central and western India are common. Thus, a research attempt to elucidate



Fig. 1. Location of pedons (P1 and P2) in the study area.

the genesis of these Mollisols is warranted. Accordingly, the present study was attempted to find out the probable reasons for the formation and persistence of Mollisols in the humid tropical climate of India. It is hoped that this addition of knowledge may help in expanding the understanding of the factors of formation of Mollisols.

2. Materials and methods

2.1. Materials

For the present investigation two Mollisols were selected from the representative catena consisting of Mollisols-Alfisols-Vertisols from Central and Western India (Fig. 1).

Pedon 1 was from the Satpura ranges of the Central India. This area is characterized by humid subtropical climate with mean annual rainfall of 1422 mm and cool temperature from November to February (mean minimum winter temperature of 8.9 °C). Majority of rainfall $(\sim 88\%)$ occurs during four wet months with a mean air temperature of 24.4 °C (Table 1). Pedon 1, on the Deccan basalt, occurs on the side slopes at an elevation of 850 m above mean sea level (msl). These areas are under thick forest vegetations of teak (Tectona grandis), mahua (Madhuca indica) and shrubs and grasses for centuries. The pedon has more than 150 cm thick solum with clay loam texture. The soils have COLE values of 0.1 and mollic epipedon which is 20 cm thick to qualify for Vertic Haplustoll subgroup in Soil Taxonomy (Soil Survey Staff, 2003). The morphological description of the soils is given in Table 2.

Pedon 2, also on the Deccan basalt, is located in the Western Ghats of India. This area is characterized by humid tropical climate with mean annual rainfall of 3287 mm and warm winter months of November, December, January and February (mean minimum winter temperature of 14.2 °C). Most of the rainfall $(\sim 91\%)$ occurs during four wet months with mean air temperature of 27.0 °C (Table 1). The Mollisols occur on the side slopes of the plateau at an elevation of 1150 m above msl and are under mixed forest (deciduous) vegetation. The soils have more than 200 cm thick solum with clay texture. They have patchy clay cutans between 40 and 100 cm depth. The dark coloured mollic epipedon, cracks and COLE (>0.06) justify grouping this pedon into Vertic Argiustoll (Soil Survey Staff, 2003). The morphological description of Pedon 2 is given in Table 2.

These Mollisols, although under forest, are prone to erosion due to slope factor. This may perhaps justify the thinner mollic epipedon even in soils under forest. The

Table I				
Climatic	datasets	of the	study	area

Months	Mean temperature (°C)	Rainfall (mm)
Satpura Ranges		
Pedon 1: Vertic I	Haplustoll	
January	17.4	28.7
February	19.7	15.3
March	23.9	27.6
April	28.5	13.2
May	32.8	10.0
June	31.4	131.7
July	26.7	480.4
August	26.1	419.5
September	26.2	213.7
October	24.0	70.6
November	19.0	8.3
December	16.9	2.8
Average	24.4	
Total		1421.8
Western Ghats		
Pedon 2: Vertic	Argiudoll	
January	25.1	2.3
February	25.3	1.0
March	27.0	0.5
April	28.9	5.6
May	29.8	27.9
June	27.9	803.4
July	26.7	1247.1
August	26.2	715.8
September	26.5	364.0
October	27.2	92.7
November	26.9	24.4
December	26.0	2.5
Average	27.0	-
Total		3287.2

recent modifications of mollic epipedon have been proposed for soils under eroded conditions. The eroded conditions have been explained as a pre-requisite for the cultivated soils (Olson et al., 2005).

2.2. Methods

The profiles were examined and described following the methods laid down in Soil Survey Manual (Soil Survey Division Staff, 1995) and Keys to Soil Taxonomy (Soil Survey Staff, 2003). The international pipette method was followed for the determination of particle-size distribution. Sand (2000–50 μ m), silt (50– 2 μ m) and clay (<2 and <0.2 μ m) fractions were separated from the samples after removing carbonates, organic matter and oxides of Fe and Al and then dispersing the soils according to the size segregation procedure of Jackson (1979). Available water content (AWC) was calculated using the water retained between 33 kPa and 1500 kPa of less than 2 mm size soil samples

Horizon	Depth (cm)	Elevation (m above msl) ^a	Slope (%)	Geology (parent material)	Drainage	Texture ^b	Structure ^c	Munsell colour ^d (moist)	Other features	Land use	Physiographic or topographic position
Pedon 1:	Village Gu	injhari, disti	rict Man	dla, Madhya .	Pradesh, Ir	ndia (Fine-l	oamy, smect	itic, hyperther	mic Vertic	Haplustoll)	
A1	0-6	850	15-30	Basalt	Very	cl	1f gr	5YR 2.5/1	Cracks	Forest (Tectona	Hill slope
A2	6-20			(weathered	well	cl	1f gr	5YR 2.5/1	$\sim\!0.5~cm$	grandis,	
Bw1	20-37			basalt)	drained	cl	1f sbk	5YR 2.5/2	wide	Madhuca	
Bw2	37-74					cl	2 m sbk	5YR 2.5/2		indica)	
Bw3	74-106					cl	2 m sbk	5YR 4/3			
Bw4	106-150					cl	2 m sbk	5YR 4/3			
Pedon 2:	Village Nig	gdale, distri	ct Pune,	Maharashtra	, India (Vei	ry fine, mix	ed, isohyper	thermic Vertic	Argiudoll)		
A1	0-15	1150	~ 40	Basalt	Very	c	1 f gr	7.5YR 3/2	Cracks	Forest	Hill slope
Bw	15 - 40			(weathered	well	с	1 f gr	7.5 YR 3/2	$\sim 0.5 \text{ cm}$	(Syzgium	
				basalt)	drained				wide	cuminii,	
Bt1	40-74					с	2 m sbk	5 YR 3/3	Clay	Terminalia	
									cutans	chebula,	
Bt2	74-108					с	3 c sbk	2.5YR 3/4	Clay	Carissa	
									cutans	caranadas,	
Bt3	108-146					с	2 m sbk	2.5YR 3/4	Clay	Ficus	
									cutans	glomerata)	
BC1	146-175					с	2 m sbk	2.5YR 3/4			
BC2	175-190					с	2 m sbk	2.5 YR 3/4			

Table 2 Morphological characteristics of Mollisols

^a msl=mean sea level.

^b cl=clay loam; c=clay.

^c 1f gr=weak fine granular; 1f sbk=weak fine subangular blocky; 2 m sbk=moderate medium subangular blocky; 3c sbk=strong coarse subangular blocky.

^d 5YR 2.5/1=Black; 5YR 2.5/2=Dark reddish brown; 5YR 4/3=Reddish brown; 7.5YR 3/2=Dark brown; 5YR 3/3=Dark reddish brown; 2.5YR 3/4=Dark reddish brown (Also see Soil Survey Division Staff, 1995).

(Richards, 1954). Extractable bases (Ca²⁺, Mg²⁺, Na⁺ and K⁺) were determined following the standard method (Jackson, 1973). Cation exchange capacity (CEC) was determined by saturating soils with 1 N NaOAc (pH 7) and leaching the Na-saturated soils by $Mg(NO_3)_2$ to eliminate the influence of Na⁺ ion from zeolites and feldspathoid minerals (Gupta et al., 1985; Pal et al., 1994). The Na⁺ ions were measured in atomic absorption spectrophotometer to calculate the CEC. Soil organic carbon was determined following the procedure of Walkley and Black (1934). The coefficient of linear extensibility (COLE) was determined following the method of Schafer and Singer (1976). The silt and clay fractions were subjected to examination of Xray diffraction (XRD) analyses of parallel oriented slide mounts after Mg- and K-saturation, Mg-glycolation using ethylene glycol, heat treatment of K-saturated samples at 25, 110, 300 and 550 °C, and HCl treatment (Jackson, 1979) using a Philips diffractometer and Nifiltered CuK α radiation at a scanning speed of 1°2 θ / min. The powdered sand samples were also examined by XRD after box mounting. Minerals were identified following various diagnostic methods (Jackson, 1979;

Brown, 1984). The semi-quantitative estimates of clay minerals in the clay fractions were carried out following the method of Gjems (1967).

3. Results

3.1. Physical and chemical properties

The Mollisol of the Western Ghats (Pedon 2) contains less amount of sand (10-16%) than Pedon 1 (23-52%), whereas total clay content was higher (51-61%) throughout the depth of Pedon 2 than Pedon 1 (28-39%). Pedon 1 has also more fine clay in the clay fractions (46-76%) than Pedon 2 (39-59%). Higher proportion of fine clay fractions as well as higher content of smectites indicates more available reactive surface of Pedon 1 per unit weight of clay than Pedon 2. Pedon 2 had Bt horizon which was not identified in Pedon 1. Both the soils have a considerable amount of AWC for biological activities and litter decomposition (Table 3).

Both the pedons have acidic reaction; however, Mollisol (Pedon 1) of the Satpura are marginally more

Horizon Depth (cm)	Depth	Particle-size distrib		Textural ^a	Available		
	(cm)	Sand (2000–50 μm)	Silt (50–2 μm)	Clay (<2 μm)	Fine clay (<0.2 μm)	class (CS)	water content (%)
Satpura Ran	iges						
Pedon 1: Ve	rtic Haplustoll						
A1	0-6	31	39	30	22 (73) ^b	Clay loam	15
A2	6-20	23	38	39	19 (46)		16
Bw1	20-37	30	41	29	22 (76)		18
Bw2	37-74	38	31	31	22 (71)		18
Bw3	74-106	42	27	31	20 (64)		19
Bw4	106-150	52	20	28	16 (57)		19
CS ^a	20-150	38	31	31	=		-
Western Gha	ats						
Pedon 2: Ve	rtic Argiudoll						
A1	0-15	16	33	51	29 (57)	Clay	15
Bw	15-40	12	35	53	31 (58)	•	17
Bt1	40-74	10	29	61	36 (59)		18
Bt2	74-108	11	28	61	35 (57)		17
Bt3	108-146	13	34	59	31 (51)		18
BC1	146-175	13	36	53	25 (47)		18
BC2	175-190	13	34	51	20 (39)		15
CS ^a	40-146	11	30	59	_	_	

Table 3 Particle-size distribution, textural class and AWC of Mollisols

^a CS : Control section; CS is defined by a depth of 150 cm from the soil surface, since the bottom of the deepest diagnostic horizon (cambic for Pedon 1: thickness 150–20=130 cm and argillic for Pedon 2: thickness 146–40=106 cm) is less than 150 cm (Soil Survey Staff, 2003). ^b Parentheses () indicate % of fine clay.

Table 4		
Physical and chemical	properties of the Mollisols	

Horizon Depth		рН (1:2)		OC ^a	Extract	able bases	(cmol (+)	kg^{-1})	CEC ^b	BS	COLEd
(cm)	Water	KCl	$(g kg^{-1})$	Са	Mg	Na	K	$(\text{cmol} (+) \text{ kg}^{-1})$	(%) ^c		
Satpura R	anges										
Pedon 1: J	Vertic Haplustol	1									
A1	0-6	5.9	5.1	35	29.6	14.4	1.0	2.3	52.2	91	0.11
A2	6-20	5.8	4.8	30	32.8	16.4	1.0	2.0	59.8	87	0.12
Bw1	20-37	5.8	4.6	20	33.2	18.8	1.1	1.8	59.8	92	0.11
Bw2	37-74	5.9	4.3	12	36.8	19.2	1.1	1.4	67.4	87	0.14
Bw3	74-106	5.6	4.2	8	45.4	18.4	1.2	0.8	71.7	92	0.16
Bw4	106-150	5.5	4.0	5	45.2	17.6	1.1	1.0	73.9	88	0.16
Western G	hats										
Pedon 2: J	Vertic Argiudoll										
A1	0-15	5.7	5.0	20	9.2	2.5	0.4	0.5	18.6	68	0.10
Bw	15-40	5.7	5.1	12	10.3	2.7	0.4	0.4	18.5	75	0.14
Bt1	40-74	5.7	5.2	7	10.3	2.8	0.4	0.2	18.7	73	0.16
Bt2	74-108	6.1	5.3	4	12.0	3.9	0.4	0.2	18.6	89	0.17
Bt3	108-146	6.1	5.0	3	12.0	4.6	0.5	0.2	18.7	92	0.15
BC1	146-175	6.1	5.0	1	11.9	5.7	0.4	0.2	20.0	91	0.13
BC2	175-190	6.1	5.4	1	11.7	5.9	0.5	0.3	19.5	94	0.13

^a OC=organic carbon.

^b CEC=cation exchange capacity.

^c BS=base saturation.

^d COLE=coefficient of linear extensibility.

acidic than that (Pedon 2) of the Western Ghats. Organic carbon content is more in Pedon 1 than Pedon 2. The CEC of soils on the Satpura (Pedon 1) was higher than those of the Western Ghats (Table 4). Higher CEC of Pedon 1 is due to more amounts of smectites. It is interesting to note that despite their acidic reaction and non-calcareous nature both the Mollisols have very high base saturation (Table 4). This is due to the presence of zeolites as confirmed by XRD studies.

3.2. Mineralogy of sand-size fractions

The sand fractions $(2000-50 \ \mu\text{m})$ of Pedon 1 contained quartz, feldspars and zeolites. The XRD analysis indicated a strong peak at 0.90 nm of zeolites. Various thermal treatments indicated that the zeolite peak at 0.90 nm remained unchanged up to 300 °C and it disappeared at 450 °C (Fig. 2). This suggests that these



Fig. 2. X-ray diffractograms of sand fractions in Mollisols as representative in terms of behaviour of zeolite mineral towards thermal treatments: (a) room temperature; (b) 210 °C; (c) 300 °C; (d) 450 °C; (e) 550 °C. Z, Zeolite; F, Feldspars; Q, Quartz.

zeolites belong to Si-poor heulandite type (Brown, 1984; Ming and Dixon, 1986; Bhattacharyya et al., 1993, 1999). The sand fractions of Pedon 2 showed a strong peak at 0.90 nm of zeolite with other accessory minerals such as mica, feldspars, quartz and anatase (figure not shown). The zeolite present in both the Mollisols is similar (Fig. 2).

The silt fractions (50–2 μ m) of both the pedons contained smectite, mica, smectite-kaolin interstratified (Sm/K) minerals, quartz and feldspars (Fig. 3). The Mollisol on the Satpura (Pedon 1) contained zeolites also. The chloritisation of smectites was indicated by incomplete expansion of the 1.40 nm peak on glycolation and broadening on the low angle side of the 1.0 nm peak of K-saturated sample heated to 550 °C. The 0.90 nm peak of zeolite disappeared at 550 °C indicating Si-poor heulandite type of zeolites. A slight shift and tailing of the 0.7 nm peak on glycolation and gradual reinforcement of the 1.0 nm peak with a corresponding decrease in the 0.7 nm peak intensity on K-saturation and subsequent heating (110-550 °C) suggested that these kaolins are to some extent interstratified with chloritised smectite (Sm/K). The formation of Sm/K is generally associated in soils developed in humid tropical climate, where it forms an important ephemeral stage during the transformation of smectite to kaolin (Pal et al., 1989; Bhattacharyya et al., 1993).

The coarse clay fractions $(2-0.2 \ \mu\text{m})$ of both the pedons were similar in mineral suit. The minerals identified were smectite, mica, Sm/K, quartz and feldspars. Zeolites were also identified in Mollisol of the Satpura ranges. These minerals belonged to Si-poor heulandite type. The Sm/K minerals were dominant in Pedon 2. Peak shift analysis (Wilson, 1987) indicates considerable amount of smectites in both the pedons (Table 5).

The fine clay fractions ($< 0.2 \mu m$) of soils contain smectites and Sm/K in different proportions. Representative XRDs are shown in Fig. 4. The identified mineral at 0.7 nm was not discrete kaolinite but are interstratified minerals of Sm/K. The relative proportion of Sm in these Sm/K interstratified minerals (Table 6) indicates that although the 0.7 nm reflection appears to be a peak of kaolinite (Hajek, 1985; Yerima et al., 1985, 1987) it actually contains little smectite (Bhattacharyya et al., 1997). This type of smectite is primarily responsible for retention of high amount of organic carbon in the Mollisols under study. Total amount of smectite including the smectites of Sm/K after the peak shift analysis (Wilson, 1987) is higher in Pedon 1 (Satpura) than Pedon 2 (Western Ghats) (Table 6). This suggests that the surface area available in Mollisol of Satpura is more than that of



Fig. 3. X-ray diffractograms of representative silt fractions of (a) P1, Bw1 (20–37 cm) and (b) P2, Bt1 (40–74 cm); Mg, Mg saturated; MgEG, Mg saturated and glycolated; K25/110/300/550 °C, K saturated and heated to 25 °C, 110 °C, 300 °C and 550 °C; 6 N HCl, samples boiled in 6 N HCl for 30 min. Sm, Smectite; M, Mica; Z, Zeolite; Sm/K, Smectite–kaolin, Q, Quartz, F, Feldspars, C, Cristobalite, A, Anatase.

the Western Ghats. This has resulted in more soil organic carbon (SOC) in former than the latter (Table 4).

4. Discussion

4.1. Formation of Mollisols

Mollisols are formed abundantly in temperate humid climate, which is conducive for the formation and retention of high organic matter (OM). Mollisols are, by and large, neutral in reaction with more than 50% base saturation and 0.6% SOC and exhibit lower degree of weathering. Some Mollisols are formed in areas where precipitation is sufficient for water to leach the soluble soil constituents and weathering products through the profile (Buol et al., 1973). Other Mollisols are formed in areas where water enters the soil and is held by the soil particles till it is lost by evapotranspiration. Percolation through the profile frequently may not occur in such areas. The soluble constituents and weathering products may not be lost

from these soils. Thus, in the environment where Mollisols are formed may have a wide range of leaching environments, stages of weathering and horizon differentiation. These ranges may include soils that have free CaCO₃ in the surface horizon to those soils in which the parent material has been leached to a depth of more than 2 m (Fenton, 1983). The process of Mollisol formation has been related to the general pedogenic process of calcification leading to the formation of chernozemic soils (Marbut, 1951; Fanning and Fanning, 1989). Calcification indicates weak eluviation and the formation of secondary calcium carbonate in sub-soil horizons. Due to the process of calcification influenced by such conditions, a stable, black type of humified organic matter forms throughout a typically thick A horizon and often the upper part of the B horizon. The mollic epipedon was thus defined to recognize the dark, high base status zone at the top of mineral soils. The dark colour of A and sometimes the upper B horizon of Mollisols may result largely due to Ca-humate which refers to Ca-

Table 5	
Semi-quantitative estimates of minerals in coarse clay fractions (2–0.2 μ m) of Mollisols	

Horizon	Depth	Minerals (%)							
	(cm)	Smectite (Sm) ^a	Mica	Zeolite	Sm/K (Sm:K) (K) ^b	Quartz	Feldspars		
Satpura Ran	nges								
Pedon 1: Ver	rtic Haplustoll								
A1	0-6	22 (32)	9	16	37 (26:74) (27)	14	Trace		
A2	6-20	23 (31)	12	22	30 (26:74) (22)	9	Trace		
Bw1	20-37	27 (36)	Trace ^c	17	36 (26:74) (27)	8	Trace		
Bw2	37-74	20 (31)	6	16	41 (26:74) (30)	15	Trace		
Bw3	74-106	32 (38)	6	18	34 (18:82) (28)	8	Trace		
Bw4	106-150	32 (38)	6	18	34 (18:82) (28)	8	Trace		
Western Gha	ats								
Pedon 2: Ver	rtic Argiustoll								
A1	0-15	13 (29)	9	Nil	62 (26:74) (46)	9	7		
Bw	15-40	12 (24)	9	Nil	68 (18:82) (56)	7	Trace		
Bt1	40-74	11 (20)	21	Nil	59 (15:85) (50)	8	Trace		
Bt2	74-108	18 (25)	9	Nil	61 (12:88) (54)	7	Trace		
Bt3	108-146	18 (26)	9	Nil	65 (12:88) (57)	6	Trace		
BC1	146-175	18 (23)	10	Nil	62 (9:91) (57)	10	Nil		
BC2	175-190	11 (17)	Trace	Nil	69 (9:91) (63)	9	9		

^a (Sm)=Effective smectite (after peak shift analysis)=% Smectite +% smectite from Sm/K [as an example say for Bw, 15–40 cm horizon of Pedon 2 effective smectite]=12%+18% of 68=(12+12)%=24%.

^b (K)=Effective kaolinite (K) content (after peak shift analysis)=82% of 68=56% (for 15-40 cm horizon of Pedon 2).

^c Trace=<5%.

saturated humic acid in the humified organic matter. Ca-ions apparently slow the rate of decomposition of Ca-humate enabling organic matter content of the soils quite high (Fanning and Fanning, 1989). This has been reported due to the formation of bridge-linked compound between inorganic clay colloids and humic acid in presence of polyvalent cations like Ca and Mg (Varadachari et al., 1991; Bhattacharyya and Ghosh, 1994).

In contrast to commonly reported calcareous Mollisols (Fanning and Fanning, 1989), the Mollisols under study are acidic and non-calcareous. Their occurrence in high weathering and leaching environments clearly points out a fact that the parent material must be providing enough of Ca ions and moisture for the formation and persistence of Mollisols.

The presence of Sm/K clay minerals indicates the transformation of smectites to kaolin in tropical humid climate of the study area (Pal et al., 1989). The clay fraction is dominated by smectite and Sm/K in both Pedon 1 and Pedon 2 and the content of smectite in Pedon 1 is more than Sm/K in Pedon 2 (Table 6). The subdominant amount of smectite holds enough moisture (Table 4) to maintain a pedo-environment for the accumulation of organic matter and has made the formation of mollic epipedon possible even in tropical humid climate. This has been possible due to zeolites, which provided continuous supply of bases not only to

arrest the complete transformation of smectite to kaolin but also to maintain very high base saturation level of the Mollisols.

Influence of the quality of parent materials in terms of clay minerals and bases in maintaining the slowly oxidizable pool, very slowly oxidizable pool and the passive or recalcitrant pool of OM has been assessed (Eswaran et al., 1995; Batjes, 2001) and the contribution of mineralogy in sequestration of OM has been recalculated to be as high as 78% (Bhattacharyya and Pal, 2003). Since mollic epipedon represents soils containing high OM content, it is expected that formation of Mollisols should be possible in forest areas with high rainfall and cool climate. The effect of vegetation and climate thus greatly influence the formation of these soils. Among the climatic parameters, rainfall favouring leaching has a dominant influence on soil properties in setting the level of base saturation. And if the rate of weathering of parent materials is high due to high rainfall and relatively warm tropical temperature, the base saturation assumes more importance as realized in Mollisols of the Western Ghats. Reports of Alfisols and Ultisols in high rainfall and cool temperature regions of thickly vegetated Indian Himalaya on poorbase rock systems (Bhattacharyya et al., 1994, 1996; Navak et al., 1996; Sidhu et al., 1997; Rana et al., 2000) clearly indicate that more rainfall and vegetation with cool temperature do not favour the formation of



Fig. 4. X-ray diffractograms of representative fine clay fractions of soils: (a) P1, Bw1 (20–37 cm) and (b) P2, Bt1 (40–74 cm); Mg, Mg saturated; MgEG, Mg saturated and glycolated; K25/110/300/550 °C, K saturated and heated to 25 °C, 110 °C, 300 °C and 550 °C; 6 N HCl, samples boiled in 6 N HCl for 30 min. Sm, Smectite; M, Mica; Sm/K, Smectite–kaolin.

Mollisols even when bases are added in soils through forest biocycling. This reaffirms the role of the quality of parent material in terms of clay minerals and bases not only in accumulation of SOC but also in the formation and persistence of Mollisols.

4.2. Persistence of Mollisols

Mollisols are abundant in temperate climate with extremely cool temperature in a landscape ranging from mid-Pleistocene to Holocene with less degree of soil weathering. The Mollisols described in the present study represent the humid tropical climate. The presence of zeolites in the coarse clay fractions (2– 0.2 μ m) indicates a huge source of base-rich heulandites, which still persist. The loss of bases during the leaching of soils has thus been continuously replenished by the steady supply of bases from heulandites. This process develops a chemical environment which prevents the formation of kaolinitic and/or oxidic clay minerals, characteristic of typical advanced stage of soil weathering. In fact these zeolitic minerals, acting as saviours against further soil degradation (Bhattacharyya et al., 2000), maintain a base saturation level (>50%) sufficiently high enough to keep these soils moist, soft and dark even with sub-dominating proportion of smectites in the present day humid tropical climate.

The Mollisol of each of the study area is a member of Mollisol-Alfisol-Vertisol association. The Mollisols of the Western Ghats have argillic horizons unlike those of the Satpura Range. This indicates that the Mollisols and the associated ferruginous Alfisols are formed in a stable (basaltic) landscape. It has been observed that the Alfisols are more acidic, have more clay and kaolinite than the Mollisols (Table 7). This suggests that the Alfisols (under sparse forest) are more weathered than Mollisols due to better water influx in the former. This indicates that even under similar parent material and climate two different groups of soils can be formed due to difference in density of forest vegetation. The associated ferruginous Alfisols were formed in tropical humid climate and are persisting since the early Tertiary (Bhattacharyya et al., 1999). The transformation of smectite to Sm/K during humid tropical weathering began at the end of the Cretaceous and continued during the Tertiary (Kumar, 1986; Tardy et al., 1991). Many of these ferruginous soils date back to the Tertiary and Cretaceous (Idnurm and Schmidt, 1986). In other words, they were formed almost as soon as the Deccan basalt was erupted (Ollier, 1995). This suggests that although the soils in the study area are formed in humid tropical climate for millions of years they have not reached the

Table 6

Semi-quantitative estimates of minerals in fine clay fractions ($<0.2 \ \mu m$) of Mollisols

Horizon	Depth	Minerals (%)					
	(cm)	Smectite (Sm) ^a	Mica	Sm/K (Sm:K) (K) ^t			
Satpura 1	Ranges						
Pedon 1 :	Vertic Hap	olustoll					
A1	0-6	68 (74)	8	24 (23:77) (18)			
A2	6-20	73 (79)	10	17 (33:67) (11)			
Bw1	20-37	76 (82)	9	15 (33:67) (9)			
Bw2	37-74	76 (85)	Trace ^c	20(43:57) (11)			
Bw3	74-106	72 (82)	Trace	23 (43:57) (13)			
Bw4	106-150	81 (93)	Nil	19 (65:35) (7)			
Western (Ghats						
Pedon 2	Vertic Arg	iustoll					
A1	0-15	12 (29)	14	74 (23:77) (57)			
Bw	15 - 40	12 (23)	14	74 (15:85) (63)			
Bt1	40-74	16 (26)	15	58 (15:85) (59)			
Bt2	74-108	12 (33)	19	69 (31:69) (48)			
Bt3	108-146	Nil (21)	8	92 (23:77) (71)			
BC1	146-175	Nil (7)	8	92 (8:92) (85)			
BC2	175-190	Nil (7)	Trace	95 (8:92) (88)			

^a (Sm)=Effective smectite (after peak shift analysis)=% Smectite+% smectite from Sm/K [As an example say for Bw, 15–40 cm horizon of Pedon 2 effective smectite]=12%+15% of 74=(12+11)%=23%.

```
<sup>b</sup> (K)=Effective kaolinite (K) content (after peak shift analysis)
=85\% of 74=63% (for 15-40 cm horizon of Pedon 2).
```

^c Trace=<5%.

Table 7	
Salient properties of Mollisols and associated Alfisols (0-30 cm	ı)

Soils	Properties	Properties							
	pH (water)	Clay (%)	Minerals (% clay fraction	Minerals (%) in the fine clay fractions					
			Smectite ^a	Kaolinite ^a					
Satpura Ra	inges								
Mollisol	5.8	33.8	79	12 (Mica 9%)					
Alfisol	5.2	50.5	37	54 (Mica 9%)					
Western Gl	hats								
Mollisol	5.7	52.0	26	60 (Mica 14%)					
Alfisol	5.4	56.0	18	67 (Mica 15%)					

^a Values after peak shift analysis (Wilson, 1987).

advanced stage of weathering represented by Ultisols and Oxisols (Soil Survey Staff, 1999). Instead they represent Alfisols and Mollisols.

The formation and persistence of the Mollisols, therefore, support that the steady state may exist in soils developed over long periods of time not only a few hundreds to thousands of years (Yaalon, 1971, 1975; Smeck et al., 1983) but also millions of years. The hypothesis of Chesworth (Chesworth, 1973, 1980) for soil formation in humid tropical climate cannot explain the persistence of these Mollisols because the stability of zeolites over time was not considered in his model. Thus the formation and persistence of these Mollisols provide an unique example that in an open system such as soil, the existence of a steady state appears to be a more meaningful concept than equilibrium in a rigorous thermodynamic sense (Smeck et al., 1983; Bhattacharyya et al., 1999). Due to the presence of zeolites the adverse effect of tropical weathering was overcome and thus the zeolitic Deccan basalt could produce a group of organic matter-rich, dark-coloured, soft, clayey, smectitic but acidic Mollisols.

5. Conclusions

The present study provides a unique example of formation and persistence of Mollisols in humid tropical environment. The Mollisols have been formed due to the conditions of better water storage effecting retention of more organic matter in the soil system to maintain the mollic epipedon. Better moisture storage has been due to the presence of smectites having more surface area. The continuous supply of bases from zeolites of amygdoloidal basalts are responsible for the stabilization of smectites in such an environment. The present model on the formation and persistence of Mollisols in the hills of central and western India expands the basic knowledge on the formation of Mollisols in humid tropical environments.

References

- Batjes, N.H., 2001. Options for increasing carbon sequestration in West African soils: an exploratory study with special focus on Senegal. Land Degrad. Dev. 12, 131–142.
- Bhattacharyya, T., Ghosh, S.K., 1994. Nature and characteristics of naturally occurring clay-organic complex of two soils from northeastern region. Clay Res. 13, 1–9.
- Bhattacharyya, T., Pal, D.K., 2003. Carbon sequestration in soils of the Indo-Gangetic Plain. Addressing Resource Conservation Issues in Rice-Wheat Systems of South Asia—A Resource Book, Rice– Wheat Consortium for Indo-Gangetic Plains. CIMMYT-Centre, New Delhi, pp. 68–71.
- Bhattacharyya, T., Pal, D.K., Deshpande, S.B., 1993. Genesis and transformation of minerals in the formation of red (Alfisols) and black (Inceptisols and Vertisols) soils on Deccan basalt. J. Soil Sci. 44, 159–171.
- Bhattacharyya, T., Sen, T.K., Singh, Nayak, D.C., Sehgal, J.L., 1994. Morphological characteristics and classification of Ultisols with kandic horizon in North Eastern Region. J. Indian Soc. Soil Sci. 42, 301–306.
- Bhattacharyya, T., Sehgal, J., Sarkar, D., 1996. Soils of Tripura for optimizing land use: their kinds, distribution and suitability for major field crops and rubber. NBSS Publ. 65b. Soils of India Series, vol. 6. NBSS&LUP, Nagpur, India, p. 154.
- Bhattacharyya, T., Pal, D.K., Deshpande, S.B., 1997. On kaolinitic and mixed mineralogy classes of shrink-swell soils. Aust. J. Soil Res. 35, 1245–1252.
- Bhattacharyya, T., Pal, D.K., Srivastava, P., 1999. Role of zeolites in persistence of high altitude ferruginous Alfisols of the Western Ghats, India. Geoderma 90, 263–276.
- Bhattacharyya, T., Pal, D.K., Srivastava, P., Velayutham, M., 2000. Natural zeolites as saviour against soil degradation. Gondwana Geol. Soc. Mag. 16, 27–29.
- Brown, G., 1984. Associated minerals. In: Brindley, G.W., Brown, G.
 W. (Eds.), Crystal Structures of Clay Minerals and Their X-ray Identification Mineralogical Society, London, pp. 361–410.
- Buol, S.W., Hole, F.D., McCracken, R.J., 1973. Soil Genesis and Classification. The Iowa State University Press, Ames, IA.
- Chesworth, W., 1973. The parent rock effect in the genesis of soil. Geoderma 10, 215–225.
- Chesworth, W., 1980. The haplosoil system. Am. J. Sci. 280, 969-985.
- Das, T.H., Thampi, C.J., Sehgal, J., Velayutham, M., 1996. Soils of sikkim for optimising land use. NBSS Publication 60b. Soils of India Series. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 44.
- Deshpande, S.B., Fehrenbacher, J.B., Ray, B.W., 1971. Mollisols of tarai region of Uttar Pradesh, Northern India: 2. Genesis and Classification. Geoderma 6, 195–201.
- Eswaran, H., van der Berg, E., Reich, P., Kimble, J., 1995. Global soil resources. In: Lal, R., Kimble, J., Levine, E., Stewart, B.A. (Eds.), Soils and Global Change. Lewis Publishers, Boca Raton, FL, pp. 27–43.
- Fanning, D.S., Fanning, M.C.B., 1989. Soil: Morphology, Genesis and Classification. John Wiley and Sons, USA, p. 395.
- Fenton, T.E., 1983. Mollisols. In: Wilding, L.P., Smeck, N.E., Hall, G. F. (Eds.), Pedogenesis and Soil Taxonomy: II. Soil Orders. Elsevier, Amsterdam, pp. 125–163.

- Gjems, O., 1967. Studies on clay minerals and clay mineral formation of soil profiles in Scandinavia. Medd. Nor. Skogforsoksves. 21, 303–415.
- Gupta, R.K., Singh, C.P., Abrol, I.P., 1985. Determining cation exchange capacity and exchangeable sodium in alkali soils. Soil Sci. 139, 326–332.
- Hajek, B.F., 1985. Mineralogy of Aridisols and Vertisols. Proceedings 5th International Soil Classification. Soil Survey Administration, Khartoum, pp. 221–230. Sudan.
- Idnurm, M., Schmidt, P.W., 1986. Paleo-magnetic dating of weathered profile. Geol. Surv. India Mem. 120, 79–88.
- Jackson, M.L., 1973. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi, India, p. 498.
- Jackson, M.L., 1979. Soil Chemical Analysis, Advanced Course, 2nd edn. University of Wisconsin, Madison, WI, USA, 11th Printing, Published by the author, pp. 895.
- Krishnan, P., Venugopal, K.R., Sehgal, J., 1996. Soil resources of Kerala for land use planning. NBSS Publication 48b. Soils of India Series, vol. 10. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 54.
- Kumar, A., 1986. Palaeolatitudes and the age of Indian laterites. Palaeogeogr. Palaeoclimatol. Palaeoecol. 53, 231–237.
- Marbut, C.F., 1951. Soils: Their Genesis and Classification. Soil Science Society of America, Madison, Wisconsin. This is a reprint of the first 10 lectures given by Dr. Marbut at the U.S. Department of Agriculture in Washington, D.C. in 1928 (quoted from Fanning and Fanning, 1989).
- Ming, D.W., Dixon, J.B., 1986. Clinoptilolite in south Texas soils. Soil Sci. Soc. Am. J. 50, 1618–1622.
- Murthy, R.S., Hirekerur, L.R., Deshpande, S.B., Venkat Rao, G.V. (Eds.), 1982. Benchmark soils of India. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 374.
- Nayak, D.C., Chamuah, G.S., Maji, A.K., Sehgal, J., Velayutham, M., 1996. Soils of Arunachal Pradesh for optimizing land use. NBSS Publ. 55b. Soils of India Series. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 54.
- Ollier, C.D., 1995. New concepts of laterite formation. In: Wadia, S., Korisettar, R. (Eds.), Quaternary Environments and Geoarchaeology of India. Memoir-Geological Society of India, vol. 32, pp. 309–323.
- Olson, K.R., Fenton, T.E., Smeck, N.E., Hammer, M.D., Ransom, C. W., Zanner, C.W., McLeese, R., Sucik, M.T., 2005. Proposed modifications of mollic epipedon thickness criteria for eroded conditions and potential impacts on existing soil classification. Soil Surv. Horiz. 46, 39–47.
- Pal, D.K., Deshpande, S.B., Venugopal, K.R., Kalbande, A.R., 1989. Formation of di- and trioctahedral smectite as an evidence for paleoclimatic changes in southern and central Peninsular India. Geoderma 45, 175–184.
- Pal, D.K., Kalbande, A.R., Deshpande, S.B., Sehgal, J.L., 1994. Evidence of clay illuviation in sodic soils of the Indo-Gangetic Plain since the Holocene. Soil Sci. 158, 465–473.
- Rana, K.P.C., Walia, C.S., Sidhu, G.S., Singh, S.P., Velayutham, M., Sehgal, J., 2000. Soils of Jammu and Kashmir for optimizing land use. NBSS Publ. 62. . Soils of India Series. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 71.
- Richards, L.A. (Ed.), 1954. Diagnosis and Improvement of Saline and Alkali Soils. USDA Agricultural Handbook, vol. 60. US Government Printing Office, Washington, DC.
- Schafer, W.M., Singer, M.J., 1976. A new method for measuring shrink-swell potential using soil paste. Soil Sci. Soc. Am. J. 40, 805–806.

- Shivaprasad, C.R., Reddy, R.S., Sehgal, J.L., Velayutham, M., 1998. Soils of Karnataka for optimizing land use. NBSS Publication 47. . Soils of India Series. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 110.
- Sidhu, G.S., Rana, K.P.C., Sehgal, J., Velayutham, M., 1997. Soils of Himachal Pradesh for optimizing land use. NBSS Publ. 57b. . Soils of India Series. National Bureau of Soil Survey and Land Use Planning, Nagpur, India, p. 44.
- Smeck, N.E., Runge, E.C.A., Mackintosh, E.E., 1983. Dynamics and genetic modelling of soil system. In: Wilding, L.P., Smeck, N.E., Hall, G.F. (Eds.), Pedogenesis and Soil Taxonomy: 1. Concepts and Interactions. Elsevier, New York, pp. 51–81.
- Smith, G.D., 1983. Historical Development of Soil Taxonomy background. In: Wilding, L.P., Smeck, L.E., Hall, G.F. (Eds.), Pedogenesis and Soil Taxonomy: I. Concepts and Interaction. Developments in Soil Science, vol. 11A. Elsevier, Amsterdam, pp. 23–49.
- Soil Survey Division Staff, 1995. Soil Survey Manual, USDA Agril. Handbook No. 18, New Revised Edition. Scientific Publishers, Jodhpur, India.
- Soil Survey Staff, 1999. Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys. USDA-SCS Agricultural Handbook 436, Second Edition. U.S. Govt. Printing Office, Washington, DC.
- Soil Survey Staff, 2003. Keys to Soil Taxonomy, 9th edition. United States Department of Agriculture, Natural Resources Conservation Service, Washington, DC.

- Tardy, Y., Kobilsek, B., Paquet, H., 1991. Mineralogical composition and geographical distribution of African and Brazilian periatlantic laterites. The influence of continental drift and tropical paleoclimates during the last 150 million years and implications for India and Australia. J. Afr. Earth Sci. 12, 283–285.
- Varadachari, C., Mondal, A.H., Ghosh, K., 1991. Some aspects of clay-humus complexation, effect of exchangeable cations and lattice charges. Soil Sci. 151, 220–227.
- Walkley, A., Black, I.A., 1934. An examination of the Degtareff method for determining soil organic matter and proposal modification of the chromic acid titration method. Soil Sci. 37, 29–38.
- Wilson, M.J., 1987. X-ray powder diffraction methods. In: Wilson, M. J. (Ed.), A Handbook of Determinative Methods in Clay Mineralogy. Chapman and Hall, New York, pp. 26–98.
- Yaalon, D.H., 1971. Soil-forming processes in time and space. In: Yaalon, D.H. (Ed.), Paleopedology, Origin, Nature and Dating of Paleosols. Israel Universities Press, Jerusalem, pp. 29–39.
- Yaalon, D.H., 1975. Conceptual models in pedogenesis. Can soil forming functions be solved? Geoderma 14, 189–205.
- Yerima, B.P.K., Colhoun, F.G., Senkayi, A.L., Dixon, J.B., 1985. Occurrence of interstratified kaolinite-smectite in El Salvador Vertisols. Soil Sci. Soc. Am. J. 49, 462–466.
- Yerima, B.P.K., Wilding, L.P., Calhoun, F.G., Hallmark, C.T., 1987. Volcanic ash-influenced Vertisols and associated Mollisols of El Salvador : physical, chemical and morphological properties. Soil Sci. Soc. Am. J. 51, 699–708.