

## Special Section

# Technological Interventions: Boon for Rice production in Konkan Region

A. K Shinde<sup>\*1</sup>, R. L Kunkerkar<sup>1</sup>, V. A. Thorat<sup>2</sup>, L. S Chavan<sup>1</sup>, M. Talathi<sup>3</sup>, P. Mandvakar<sup>3</sup>, A. C. Devrukhkar<sup>2</sup>, V. K. Patil<sup>4</sup>, A. D. Rane<sup>4</sup>, S. B. Dodake<sup>5</sup>, J. M. Talathi<sup>2</sup>, P. M. Haldankar<sup>6</sup> and T. Bhattacharyya<sup>6</sup>

<sup>1</sup>Regional Agricultural Research Station (DBSKKV), Karjat, Raigad, Maharashtra (India) 410201

<sup>2</sup>College of Agriculture (DBSKKV), Dapoli, Ratnagiri, Maharashtra (India) 415712

<sup>3</sup>Krishi Vigyan Kendra (DBSKKV), Roha, Raigad, Maharashtra (India) 402116

<sup>4</sup>College of Forestry (DBSKKV), Dapoli, Ratnagiri, Maharashtra (India) 415712

<sup>5</sup>Kharland Research Station (DBSKKV), Panvel, Raigad, Maharashtra (India) 410206

<sup>6</sup>Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Ratnagiri, Maharashtra (India) 415712

## Abstract

Rice is one of the premier cereal crops of the World and staple food of more than half of the World's population. In India, introduction of semi-dwarf high yielding varieties was instrumental in increasing the rice production. The steady increase in rice production over the years transformed the country from food deficit to net surplus. Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli has released and notified 25 rice varieties including 5 hybrids and has developed improved package of practices for cultivation of rice crop since 1972. The rice production has increased from 10.06 lakh tonnes (1970) to 15.69 lakh tonnes (2016) and productivity from 2.3 t·ha<sup>-1</sup> to 4.35 t·ha<sup>-1</sup>.

**Keywords:** Rice varieties, Package of practices, Doubling Farmers' Income, DFI

## Rice Scenario

Rice is staple food of India as well as Asia, Latin America, part of Africa and the Middle-East. In India, rice cultivation is taken below sea level (Kerala) up to altitude of 2000 m above MSL (Kashmir). It plays vital role in national food security and it is means of livelihood of millions of people making a slogan "Rice

is Life" most appropriate.

Globally, the highest area under rice is in India (43.86 million ha) followed by China (30.16 million ha). However, highest production of rice is in China (144.85 million tonnes) followed by India (104.80 million tonnes). This is due to higher productivity of rice in China (6.86 t ha<sup>-1</sup>) than India (3.77 t ha<sup>-1</sup>) (Table 1) (Anon. 2017)

In India, rice is cultivated in both winter and summer seasons. West Bengal, Uttar Pradesh, Andhra Pradesh, Punjab, Tamil Nadu, Bihar, Orissa, Assam, Karnataka & Haryana are 10 major rice producing States which account for more than 80 per cent rice production in India (Table 2) (Anon. 2016). More than 5000 varieties of rice are grown in India. India is world's largest exporter of Basmati rice to Saudi Arabia and other Middle-East countries, Europe and United States (Anon. 2009).

In Maharashtra, rice is cultivated on 15.56 lakh ha area in all four regions viz. Vidarbha (8.15 lakh ha), Konkan (3.69 lakh ha), Western Maharashtra (3.55 lakh ha) and Marathwada (0.156 lakh ha). The highest productivity of rough rice was in Konkan region (4.25 t ha<sup>-1</sup>) followed by Western Maharashtra (3.5 t ha<sup>-1</sup>) and Vidarbha (3.4 t ha<sup>-1</sup>) (Table 3) (Anon. 2017).

The rice production in Maharashtra in 1960 was 19.84 lakh tonnes which increased to 52.96 lakh tones in 2016 (Table 3). Thus during last 56 years the rice production has been increased by 5 times. The rice productivity in 1960 was 1.52 t ha<sup>-1</sup> and it has increased to 3.4 t ha<sup>-1</sup> in 2016 (Anon. 2017). This increase in rice production and productivity was due to cultivation of newly released

\*Corresponding author: shindeak53@rediffmail.com

Date received: 20.11.2017; Date accepted: 22.3.2018

## Doubling Farmers' Income (DFI) through DBSKKV Interventions

**Table 1.** Area, production and productivity of rice in various countries of the world (2016-17) (Anon. 2017)

| Sr. No. | Country       | Area (million ha) | Production (million tonnes) | Productivity Rough Rice (t ha <sup>-1</sup> ) |
|---------|---------------|-------------------|-----------------------------|-----------------------------------------------|
| 1.      | China         | 30.16             | 144.85                      | 6.86                                          |
| 2.      | India         | 43.86             | 104.80                      | 3.77                                          |
| 3.      | Indonesia     | 12.24             | 37.15                       | 4.78                                          |
| 4.      | Bangladesh    | 11.77             | 34.58                       | 4.41                                          |
| 5.      | Vietnam       | 7.70              | 27.86                       | 5.79                                          |
| 6.      | Thailand      | 10.08             | 18.60                       | 2.80                                          |
| 7.      | Burma         | 7.00              | 12.40                       | 2.77                                          |
| 8.      | Phillipines   | 4.60              | 11.50                       | 3.97                                          |
| 9.      | Brazil        | 2.00              | 8.16                        | 6.00                                          |
| 10.     | Japan         | 1.57              | 7.78                        | 6.81                                          |
| 11.     | United States | 1.25              | 7.12                        | 8.11                                          |
| 12.     | Pakistan      | 2.69              | 6.80                        | 3.80                                          |
|         | Total World   | 159.91            | 483.1                       | 4.50                                          |

**Table 2.** Statewise area, production and productivity of Rice in India. (Anon. 2016)

| Sr. No. | State          | Area (million ha) | Production (million tonnes) | Productivity (t ha <sup>-1</sup> ) |
|---------|----------------|-------------------|-----------------------------|------------------------------------|
| 1.      | West Bengal    | 5.39              | 14.71                       | 2.73                               |
| 2.      | Uttar Pradesh  | 5.87              | 12.22                       | 2.08                               |
| 3.      | Andhra Pradesh | 3.81              | 11.57                       | 3.04                               |
| 4.      | Punjab         | 2.89              | 11.11                       | 3.84                               |
| 5.      | Orissa         | 4.17              | 8.29                        | 1.99                               |
| 6.      | Bihar          | 3.27              | 6.38                        | 1.95                               |
| 7.      | Chattisgarh    | 3.81              | 6.02                        | 1.58                               |
| 8.      | Tamil Nadu     | 1.83              | 5.84                        | 3.19                               |
| 9.      | Assam          | 2.28              | 4.86                        | 2.13                               |
| 10.     | Haryana        | 1.29              | 4.01                        | 3.11                               |
| 11.     | Karnataka      | 1.30              | 3.66                        | 2.82                               |
| 12.     | Madhya Pradesh | 2.15              | 3.63                        | 1.69                               |
| 13.     | Jharkhand      | 1.50              | 3.32                        | 2.21                               |
| 14.     | Maharashtra    | 1.55              | 2.93                        | 1.89                               |
| 15.     | Gujarat        | 0.79              | 1.64                        | 2.08                               |
| 16.     | Kerala         | 0.20              | 0.56                        | 2.80                               |
| 17.     | Other          | 1.78              | 4.07                        | 2.29                               |
|         | All India      | 43.86             | 104.80                      | 2.44                               |

rice varieties and hybrids grown under recommended package of practices.

Konkan region of Maharashtra has 5 districts viz. Thane, Palghar, Raigad, Ratnagiri and Sindhudurg. Agroecologically the Konkan region is a part of hot-humid ecoregion consisting Sahyadri mountain and Western coastal plains extending over state of Maharashtra, Goa, Karnataka and Kerala (Sehgal *et al.* 1992). Highest productivity in Konkan is in Sindhudurg district (4.89 t ha<sup>-1</sup>) followed by Ratnagiri (4.66 t ha<sup>-1</sup>), Palghar (4.1 t ha<sup>-1</sup>), Raigad (3.99 t ha<sup>-1</sup>) and Thane (3.88 t ha<sup>-1</sup>) (Table 4).

### Technology Development in India –Historical

ICAR supported research stations have been the backbone of technology development for rice in India. These research stations collected the local rice varieties, evaluated them under their local conditions and identified the best performing varieties. Initially, to increase production and productivity, more than 445 varieties were recommended by pure-line selection for preference over the local landraces. Given the negligible use of chemical fertilizers and/or insecticides in pre-independence era, the improved varieties were mostly bred under (and were adapted to) low input conditions only.

After the establishment of the Central Rice Research Institute (CRRI) at Cuttack in 1946 by the Government of India, rice research and training received an added impetus. The traditional rice varieties already collected and maintained by the various rice research stations in the country were screened for their yield potential.

Meanwhile, the ICAR launched the All-India Coordinated Rice Improvement Project (AICRIP) in 1965 that helped in co-ordinating interdisciplinary and interinstitutional rice research within the country for improving production, productivity and profitability of rice in India. This was the beginning of moving toward self-sufficiency in rice production (Swarna *et al.* 2010).

India operated its most intensive rice breeding program under the AICRIP and achieved remarkable success. Padma and Jaya were the first two varieties that emerged from the programme. During the span of next 35 years (1965 to 2000), 632 high yielding rice varieties were released by the Central Variety Release Committee and by the State Variety Release Committees. These Varieties were bred for various ecological stress situations, or to meet the new challenges faced due to diseases and insect pests which were earlier minor but became major, or for

## Special Section

**Table 3.** Information on Area, Production and Productivity of rough rice in Maharashtra State

| Regions             | Area<br>(lakh ha) |       | Rough Rice<br>Production<br>(lakh t) |       | Rough Rice<br>Productivity<br>(t ha <sup>-1</sup> ) |      |
|---------------------|-------------------|-------|--------------------------------------|-------|-----------------------------------------------------|------|
|                     | 1960              | 2016  | 1960                                 | 2016  | 1960                                                | 2016 |
| Konkan              | 4.17              | 3.69  | 7.65                                 | 15.69 | 1.83                                                | 4.25 |
| Western Maharashtra | 2.82              | 3.55  | 4.26                                 | 12.77 | 1.51                                                | 3.50 |
| Vidarbha            | 5.21              | 8.15  | 7.08                                 | 24.46 | 1.36                                                | 3.40 |
| Marathwada          | 0.80              | 0.15  | 0.84                                 | 0.028 | 1.05                                                | 0.48 |
| Maharashtra State   | 13.00             | 15.56 | 19.84                                | 52.96 | 1.53                                                | 3.40 |

**Table 4.** Second advance Estimate of Area, Production (P) & Productivity (Py) of Kharif rice in Konkan region, Maharashtra during 2016. Area in “00” ha, P in “00” Tonnes, Py in t ha<sup>-1</sup>.

| Sr. No. | District   | Area | Rice  |       | Rough Rice |       |
|---------|------------|------|-------|-------|------------|-------|
|         |            |      | P     | Py    | P          | Py    |
| 1       | Thane      | 551  | 1476  | 2.679 | 2139       | 3.882 |
| 2       | Palghar    | 756  | 2172  | 2.873 | 3147       | 4.163 |
| 3       | Raigad     | 1094 | 3017  | 2.758 | 4372       | 3.997 |
| 4       | Ratnagiri  | 686  | 2203  | 3.214 | 3192       | 4.658 |
| 5       | Sindhudurg | 604  | 1958  | 3.242 | 2837       | 4.698 |
|         | Total      | 3690 | 10827 | 2.934 | 15691      | 4.252 |

grain quality especially the Basmati types for export (Prasad Rao 2004).

Research programme was initiated in early 1970 to develop hybrid rice in the country. However, there was no breakthrough in this program for two decades. The research program was accelerated and intensified from 1989 onwards with a mission mode project. With this concerted research effort, a remarkable success was achieved within a short span of five years and half a dozen hybrid rice varieties were developed from public and private sectors. The first four hybrid rice varieties were released in the country during 1994. By the end of 2016, a total of 76 hybrid varieties were released (Anon. 2016).

Regional Agricultural Research Station (RARS), Karjat was established by British Government in 1919 for development of rice varieties for Raigad & Thane districts. From this period RARS, Karjat has developed and released six rice varieties by Pure-line Selection

**Table 5** Rice varieties developed and released by RARS, Karjat from 1919 to 1970.

| Sr. No.                                                                | Variety (Parentage)                          | Year of release | Duration in Kharif (days) | Grain type | Yield (q ha <sup>-1</sup> ) |
|------------------------------------------------------------------------|----------------------------------------------|-----------------|---------------------------|------------|-----------------------------|
| Rice varieties developed by pure line selection method                 |                                              |                 |                           |            |                             |
| 1.                                                                     | Kolamba-42                                   | 1924            | 130-135                   | SS         | 29                          |
| 2                                                                      | Zinia-31                                     | 1934            | 125-130                   | SS         | 24                          |
| 3                                                                      | Zinia 149                                    | 1934            | 145-150                   | MS         | 29                          |
| 4                                                                      | Kada-68-1                                    | 1946            | 115-120                   | LB         | 24                          |
| 5                                                                      | Mahadi 4-4                                   | 1954            | 115-120                   | LB         | 24                          |
| 6                                                                      | Garvel 1-8                                   | 1954            | 150-152                   | LB         | 23                          |
| Rice varieties developed by hybridization followed by selection method |                                              |                 |                           |            |                             |
| 1                                                                      | Kolamba-540 (K-164 x K-26)                   | 1934            | 130-135                   | SS         | 24                          |
| 2                                                                      | Bhadas-1303 (1-B-12-11 x C-48)               | 1954            | 135-140                   | SB         | 37                          |
| 3                                                                      | Zinia 14 ((K-226 x Z-242) x (Z-149 x K-540)) | 1958            | 145-150                   | MS         | 33                          |
| 4                                                                      | Zinia-63 (Z-31 x K-540) x (Z-149)            | 1959            | 125-130                   | SS         | 25                          |

(Table 5) and four rice varieties by hybridization. These varieties have productivity of 24 to 37 q ha<sup>-1</sup>. In Maharashtra total 62 rice varieties have been released including 5 hybrids by four Agricultural Universities.


### Interventions by Dr. Balasaheb Konkan Krishi Vidyapeeth, Dapoli

#### Varietal development


After establishment of Dr. Balasaheb Konkan Krishi Vidyapeeth, Dapoli in 1972 University has released and notified 25 rice varieties including 5 hybrids. Rice varieties having Productivity 3 to 5 t ha<sup>-1</sup> and hybrids having productivity 5.5 to 7.5 t ha<sup>-1</sup>. (Table 6)

Among these rice varieties, Karjat-3 and Karjat-7 are largely accepted varieties by farmers. Karjat-3 is short duration (115-120 days), short bold grain, high yielding (4.5-5.0 t ha<sup>-1</sup>) rice variety accepted by all districts (Thane, Palghar, Raigad, Ratnagiri and Sindhudurg).

## Doubling Farmers' Income (DFI) through DBSKKV Interventions



**Fig. 1.** Change in area, production and productivity of rice in Maharashtra state and its various regions.



It is also accepted by farmers of Goa. The seed of this variety is used for *poha* making, *bhakri* and grains (food). Karjat-7 is also short duration (115-120 days), long slender grain with high yield (4.5-5.0 t ha<sup>-1</sup>) accepted by farmers. In 2016, among all the varieties whose seed was supplied in Maharashtra, Karjat-3

(4345 q) and Karjat-7 (3396 q) followed Jaya (5921 q) (Table 7).

Karjat-5 is a mid-late duration (125-130 days) variety with high yield (45-55 q ha<sup>-1</sup>) having long bold grains. This variety has been accepted in Raigad district on small scale. Karjat-8 and Karjat-9 are newly released (notified) very fine grain (short slender) rice varieties accepted by farmers. Due to short slender grain of Karjat-8 and medium slender grain type of Karjat-9, these rice varieties are fetching higher price in market. RTN-1 and RTN-24 rice varieties are very much popular and accepted by farmers of Western Maharashtra (Kolhapur, Satara) region. Ratnagiri-5 is early duration (115-120 day) short slender grain type, good yielding (36 q ha<sup>-1</sup>) notified rice variety, needs extensive extension programmes. Karjat-8 is late duration (140-145 days) short slender and Karjat-9 is midlate duration (120-125 days), medium slender rice variety. (Kunkerkar *et al.* 2014)

Among the limited available options to increase production and productivity of rice, hybrid rice is most practically feasible and readily adaptable one. University released five rice hybrids of early, midlate and late duration. Which is very popular in state as well as in country. Sahyadri is the 1<sup>st</sup> hybrid rice in Maharashtra having midlate duration (125-130 days), long slender grain type and average yield 6.5 to 7.0 t ha<sup>-1</sup>. Moderately resistant to major pests and diseases, suitable for beaten rice & parched rice. Sahyadri-4 is centrally released hybrid having early duration (115-120 days) long slender grain type. Due to its wider adaptability in varied agro-ecological conditions and earliness it was recommended and released for commercial cultivation in five states of country viz. Punjab, Haryana, Uttar Pradesh, West Bengal and Maharashtra.

### Package of practices

On various research stations of University, the research trials were conducted in various disciplines. Agronomical trials for control of weeds, spacing & fertilizer trials, rice cropping systems etc. were conducted continuously for 3 to 4 years and recommendation emerged for increasing rice yield are summarized as below. In soil science discipline, the trials on use of organic, inorganic fertilizers, slow-releasing fertilizers, rice husk etc. were conducted & recommendations were given for increasing the rice yield. For control of insect pests and diseases, various plant protection measures were studied for rice and were recommended. The integrated use of this technology by farmers results into increased in

**Special Section****Table 6.** Released and notified varieties of rice by Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli (1972 to 2016)

| Variety      | Year of release | Duration (days)            | Yield (q ha <sup>-1</sup> ) | Characteristics                                                                                                                              | Remarks                          |
|--------------|-----------------|----------------------------|-----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------|
| Ratnagiri-1  | 1986            | 110-115                    | 47-50                       | Long bold grain, mid-tall, good tillering ability dark green foliage, non-lodging, non-shattering.                                           | Notified No.SO.386-E 15.05.1990  |
| Ratnagiri-2  | 1986            | 145-155                    | 50                          | Short bold grain, mid-tall, good tillering ability, dark green foliage, non-lodging, non-shattering.                                         | Notified No.SO.386-E 15.05.1990  |
| Ratnagiri-3  | 1993            | 140-142                    | 42                          | Late duration, mid-tall, non-lodging, non-shattering, long bold grain.                                                                       | Notified No.SO.1-E 1.01.1996     |
| Ratnagiri-4  | 2009            | 125-130                    | 49                          | Long slender grain                                                                                                                           | Notified No.SO.2137-E 31.08.2010 |
| Ratnagiri-24 | 2009            | 105-115                    | 36                          | Short slender translucent grain, erect growing, compact panicle, mid tall, early variety wing light green narrow erect leaves.               | Notified No.SO.2137-E 31.08.2010 |
| Ratnagiri-73 | 1979            | Kharif-95-98<br>Rabi- 120  | 45                          | Long bold grain, dwarf, very early, non-lodging variety with erect dark green leaves.                                                        | Notified No.SO.867-E 26.11.1986  |
| Karjat-1     | 1985            | 105-110                    | 35-45                       | Short bold grain, dwarf, non lodging, dark green foliage, early duration, non-shattering responsive to N fertilizers.                        | Notified No.SO.386-E 15.05.1990  |
| Karjat-2     | 1993            | 135-140                    | 40-45                       | Long slender, fine grain, dwarf, erect with dark green leaves, flag leaf erect.                                                              | Notified No.SO.1-E 1.01.1996     |
| Karjat-3     | 1994            | 110-115                    | 45-50                       | Short bold coarse grain, early, high yielding, medium tillering, erect flag leaf with dark green colour.                                     | Notified No.SO.1-E 1.01.1996     |
| Karjat-5     | 2006            | 125-130                    | 43-53                       | Long bold grain, Resistant to blast                                                                                                          | Notified No.SO.1703-E 5.10.2007  |
| Karjat-6     | 2006            | 130-135                    | 35-40                       | Fine grain Resistant to blast                                                                                                                | Notified No.SO.1703-E 5.10.2007  |
| Karjat-7     | 2007            | Kharif 115-120<br>Rabi 130 | 45-50                       | Long slender grain                                                                                                                           | Notified No.SO.449-E 11.02.2009  |
| Karjat-184   | 2009            | 100-105                    | 30-35                       | Medium slender grain, early duration, short statured (70-75 cm height) broad dropping leaves, fine grain with blackish straw panicle colour. | Notified No.SO.2137-E 31.08.2010 |
| Phondaghat-1 | 2000            | 115-120                    | 45-50                       | Long slender grain, semi-dwarf, early, non-lodging, non-shattering.                                                                          | Notified No.SO.1177-E 25.08.2005 |
| Panvel-1     | 1984            | 125-130                    | 40-43                       | Short bold, white kernel, semi-dwarf, mid-late, faint purple tinch at older leaf sheath.                                                     | Notified No.SO.540-E 24.07.1985  |

**Doubling Farmers' Income (DFI) through DBSKKV Interventions****Table 6** continued...

|             |      |         |       |                                                                                                                         |                                       |
|-------------|------|---------|-------|-------------------------------------------------------------------------------------------------------------------------|---------------------------------------|
| Panvel-2    | 1987 | 110-115 | 33-41 | Long slender and translucent grain, semi-dwarf, non-lodging, early duration non-shattering and salt tolerant.           | Notified No.SO.386-E 15.05.1990       |
| Palghar-1   | 1988 | 120-125 | 40-45 | Medium slender grains, semi-dwarf, mid-late, high yielding.                                                             | Notified No.SO.386-E 15.05.1990       |
| Sahyadri    | 1998 | 125-130 | 60-65 | Long slender, grain, white kernel, slight aroma, mid-late, mid tall, non-lodging, non-shattering.                       | Notified No.SO.821-E 13.09.2000       |
| Sahyadri-2  | 2006 | 115-120 | 60    | Non lodging, mid tall (15 to 120 cm), midlate duration, long slender, translucent kernel.                               | Notified No.SO.122-E 6.02.2007        |
| Sahyadri-3  | 2006 | 125-130 | 65-75 | Long slender grain                                                                                                      | Notified No.SO.122-E 6.02.2007        |
| Sahyadri-4  | 2006 | 115-120 | 55-65 | Long slender grain, wide adaptability, release for five state Panjab, Haryana, Uttar pradesh, West Bengal & Maharashtra | Notified No. 454-E 20.11.2009         |
| Ratnagiri-5 | 2012 | 115-120 | 36    | Short slender grain, moderately resistant to leaf blast                                                                 | Notified S.O.2238E-23 Dated 29.6.2017 |
| Sahyadri-5  | 2012 | 140-145 | 66    | Long slender grain, moderately resistance to leaf blast, tolerant to brown plant hopper.                                | Notified S.O.2238E-24 29.6.2017       |
| Karjat-8    | 2012 | 140-145 | 35-40 | Short slender growth, moderately resistant to blast                                                                     | Notified                              |
| Karjat-9    | 2014 | 120-125 | 45-50 | Short slender grain, resistant to blast.                                                                                | Notified                              |

rice production and income.

**Cultivation practices**

- 1) In North Konkan coastal zone of Maharashtra for obtaining maximum yield and economic returns from Sahyadri rice hybrid, 25 days old seedlings be transplanted at 20 x 15 cm spacing with two seedlings per hill and be supplemented with fertilizer dose of 150:50:50 kg NPK ha<sup>-1</sup>.
- 2) To get maximum yield and net profit from Rice-Maize cropping system, rice and maize crops be fertilized with 150:75:75 kg NPK ha<sup>-1</sup> and 180:90:60 kg NPK ha<sup>-1</sup> respectively.
- 3) To get higher yields and net returns from kharif rice 48 hours *Rahu* (sprouted seeds) of rice be sown by using 8 rows DRR drum seeder on puddled field.
- 4) In direct seeded rice for control of weed oxydiarzon 0.4kg a.i ha<sup>-1</sup> or oxydiagil 100 gm a.i ha<sup>-1</sup> foliar spray be followed. First spray before imergence of weed (2-3 days after seed sowing) and second spray 45 days after

first spray be done in lateritic soil of Konkan region.

- 5) For lateritic soil of Konkan region in Groundnut and Rice cropping system foliar spray of flucloalalin 1.5 kg a.i. ha<sup>-1</sup> pre emergence weedicide be done for rabi groundnut and for rice oxydiarzon 0.5 kg a.i. ha<sup>-1</sup> sprayed before weed immergence.
- 6) For getting higher returns in well drained rice field rice seed be dibbled before monsoon & pre emergence weedicide prtilachlor (50EC) @ 0.5 kg a.i. ha<sup>-1</sup> be sprayed on wet land.
- 7) For getting higher return from direct seeded rice during kharif season general package of practices be followed. During rabi season lablabbean be grown without cultivation practices except hand weeding at 20 & 40 days of sowing or spray of pre-emergence weedicide oxydiargil 0.12 kg ha<sup>-1</sup>.
- 8) In north Konkan region for direct seeded rice pre-emergence weedicide spray oxyflurphen 300 gm ha<sup>-1</sup> & post-emergence weedicide 2-4, D 500 gm ha<sup>-1</sup> be

## Special Section

**Table 7.** Certified Seed distribution of rice in Maharashtra State (2016)

| Sr. No. | Name of Variety | Seed Sold (Q) |
|---------|-----------------|---------------|
| 1       | Jaya            | 5921          |
| 2       | Karjat-2        | 1131          |
| 3       | Karjat-3        | 4345          |
| 4       | Karjat-5        | 935           |
| 5       | Karjat-7        | 3396          |
| 6       | Masuri          | 545           |
| 7       | RTN-24          | 540           |
| 8       | RTN-1           | 619           |

sprayed or weeding be done at 20 & 40 days of sowing.

9. For kharland of Konkan region 100 kg seed be used for direct seeded kharif rice.

### Nutrient management

1. It is recommended to use Urea and Suphala (15:15:15) in 1:1.5 proportion to prepare briquettes by using briquetter machine. Further application of Urea-Suphala briquettes is recommended as alternative to Urea-DAP briquettes to rice crop increase yield and to decrease fertilizer cost.

2. It is recommended to apply Glyricidia mulch @ 1 t ha<sup>-1</sup> in between two lines along with recommended dose of fertilizers for getting healthy rice seedlings and effective weed control.

3. It is recommended to use Urea-Godawari (14:35:14) in 1.5:1 proportion to prepare briquettes by using briquetter machine. Further application of Urea-Godawari briquettes is recommended as alternative to Urea-Suphala briquettes to rice crop to increase yield and to decrease fertilizer cost.

4. To get higher yield and returns from hybrid rice (Sahyadri) under lateritic soil application of Glyricidia 5 t ha<sup>-1</sup>, N:P:K @ 150:50:75 kg ha<sup>-1</sup> be followed. Nitrogen split dose be given 50 % at transplanting 25 % in two splits at 1 month duration.

5. To get higher returns in lateritic soil in rice recommended dose of NPK be reduced upto 25% and application of FYM, poultry manure, fish meal, neem cake, organic manure be applied @ 2 t ha<sup>-1</sup>.

6. In south Konkan coastal zone for getting higher yield & returns from hybrid rice Sahyadri 1 kg m<sup>-2</sup>.

Rice burned husk be applied. Transplanting be done in double row (15x15x25 cm) & application of Glyricidia 5 t ha<sup>-1</sup> & 56 kg N+ 32 Kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and DAP briquettes be done.

7. For getting higher yield and returns from rice & reduction in cost of fertilizers the rice nursery be applied with 0.5 kg m<sup>-2</sup> rice burned husk. Rice field be applied with 2.5 t ha<sup>-1</sup>. Glyricidia leaves with Konkan Annapurna briquets 62,500 briquets ha<sup>-1</sup> instead of urea DAP briquets.

8. For getting high yield and returns from rice, Silicon based Konkan Annapurna briquettes (31:14:6:0.44 N, P, K & Silicon ) be used.

### Pest Management

1. It is recommended to apply Carbofuran 3G @ 16.5 kg ha<sup>-1</sup> for management of rice caseworm, when the pest reaches ETL and second application should be given fifteen days after first application, if necessary.

2. One spray of cartap hydrochloride 50 SP at the rate of 600 g per 500 lit of water per ha is recommended for the management of rice caseworm. Application should be given, when the pest reaches ETL (two newly infested leaves per hill).

3. Harvesting of rice by “vaibhav sickle” for the management of stem borer

4. A low cost eco friendly integrated approach involving cultivation of stem borer tolerant variety ‘Ratna’, use of 20 pheromone traps ha<sup>-1</sup> and need based application of insecticides viz. Carbofuran 16.5 kg ha<sup>-1</sup> or Cartap hydrochloride 300 g a.i. ha<sup>-1</sup> appeared to be effective in reducing stem borer infestation.

5. *Scirpophaga incertulus*, *Cnaphalocrosis medinalis* and *Leptocorisa oritorius* are the only species of yellow stem borer, leaf folder and gundhi bug, respectively, existing in the Konkan region of Maharashtra.

6. Yellow stem borer and gundhi bug appeared to be the major pests with their peak population during fourth week of September.

### Disease management

1. Three sprays of Tricyclazole (75 Wp) 10 g in 10 L of water (0.1%) or Isoprothiolane (40 EC) 10 ml in 10 L of water (0.1%) are recommended at an interval of 21 days starting from the initiation of blast symptoms for effective management of blast disease of rice.

2. Blue green algae @ 10 kg ha<sup>-1</sup> was recommended

## Doubling Farmers' Income (DFI) through DBSKKV Interventions

in paddy field of Konkan region for replacing 25 kg nitrogen.

3. Incorporation of Azolla @ 6 t ha<sup>-1</sup> was recommended in paddy field of Konkan region to save 30 kg in organic Nitrogen ha<sup>-1</sup>.

4. For effective management of brown spot of rice spray of Mancozeb @ 0.25% or Zineb @ 0.20 % or Copper Oxyclozide @ 0.25% was recommended.

5. For effective management of green algae in paddy field during Rabi season in North Konkan Coastal Zone application of 10 kg Azolla ha<sup>-1</sup> was recommended.

### *Rice economics*

1. Monocropping of Kharif rice was not profitable in Konkan region due to high labour cost, low yields, minimum use of fertilizers, manures, plant protection measures, predominance of small holding, slow acceptance of technological improvement, poor financial conditions and negative attitude of the farmers towards acceptance of crop and crop insurance policies.

2. The rice-oilseeds (groundnut), rice-vegetables, (orka, chilli, tomato, brinjal, cucumber, watermelon, onion, little gourd, bitter gourd) cropping systems are more profitable and more employment generating (480-658 days ha<sup>-1</sup> year<sup>-1</sup>) systems.

3. Greater crop diversity was observed in Thane district of North Konkan. Overall crop intensity of North Konkan (111%) is stable from last two decades.

4. There is considerable potential for crop intensification and diversification in north Konkan provided irrigation facilities and better market infrastructure for disposal of crop production. Location specific strategy of diversification towards high value crops is the necessary task.

5. The technology of drum seeder in rice cultivation must be popularized to overcome the scarcity of labour and increasing wage rates and reducing the cost of cultivation by conducting the large scale demonstrations of drum seeder (direct seeded method) by the department of Agriculture, NGOs and KVKs)

### *Integrated Farming System*

1. In North Konkan Coastal Zone of Maharashtra to get higher yields and economic returns from Kharif rice and Rabi-hot weather rice system, 100 per cent RDF through inorganics be applied to both the crops. However, to improve soil fertility and sustain productivity of Rice under 'Rice – Rice' cropping system, 50% RDF

as inorganics (50, 25, 25 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup>) be integrated with rest 50 kg N (RDN) in the form of Glyricidia green leaf manuring during Kharif and 100% RDF as inorganic (120, 50, 50 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup>) during Rabi-hot weather season.

2. In north Konkan Coastal Zone of Maharashtra, 'Rice-Brinjal' cropping system is the most profitable one followed by 'Rice-Fodder maize', 'Rice-Cabbage' and 'Rice-Groundnut' systems which may be followed by the farmers depending upon their own needs and local demand for produce in the market.

3. In North Konkan Coastal Zone of Maharashtra 'Rice - Sweet corn' cropping system be fertilized with 100 % recommended NPK as inorganics followed by application of 50 per cent RDF through inorganics alongwith 50 per cent N through FYM to secure maximum yield, net returns and B:C ratio.

4. In North Konkan Coastal Zone, 'Rice-Brinjal' cropping system is found to be most profitable followed by 'Rice-Sweet corn' cropping system which may be grown with minimum tillage by applying 125 per cent RDF to both systems.

5. For yield maximization under rice – rice cropping system of North Konkan Coastal Zone of Maharashtra, hybrid rice be fertilized @ 150:100:150:0.8:10:6 kg N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, B, Fe, Zn ha<sup>-1</sup> during Kharif and @ 150:100:150 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup> during Rabi – hot weather season.

### *Organic rice*

1. In North Konkan Coastal Zone of Maharashtra, Rice-Sweet corn system supplied with inorganic sources of nutrients followed by the same system supplied with integrated and organic nutrient management was found to be most profitable. However, to improve soil fertility and sustain productivity, Rice-Groundnut and Rice-Dolichos bean systems also be adopted with organic sources of nutrients.

2. In North Konkan Coastal Zone of Maharashtra for rice based organic cropping system, rice as Kharif crop and cucurbits viz; red pumpkin and cucumber as Rabi crops be grown. Kharif rice be manured with 1/3<sup>rd</sup> RDN through FYM, 1/3<sup>rd</sup> RDN through rice straw and 1/3<sup>rd</sup> RDN through Glyricidia green leaf manuring, whereas, Rabi cucurbits be manured as 1/3<sup>rd</sup> RDN through FYM, 1/3<sup>rd</sup> RDN through neem cake and 1/3<sup>rd</sup> RDN through vermicompost, respectively, so as to obtain higher yields and economic returns.

## Special Section

### Strategies for increasing Rice Production

In a nutshell, the all above recommendations over the time were integrated to formulate a strategy for increasing rice production and productivity in Konkan region. The highlights of this strategy are given below-

•Conservation Agriculture for increasing organic carbon content of soil (minimum tillage, stubble mulch)

•Nursery management

- 1) Ploughing, harrowing
- 2) Preparation of raised beds of proper size
- 3) Incorporation of FYM 100 kg R<sup>-1</sup>-before sowing, at sowing 3kg suphala 15:15:15 R<sup>-1</sup> and Urea 1kg R<sup>-1</sup> 15 DAS
- 4) Rice husk ash 0.5 to 1 kg m<sup>-2</sup> on nursery area
- 5) Sowing seed on raised bed (40-50 kg ha<sup>-1</sup>).
- 6) Application of Pre emergence weedicide after sowing seed & on wet bed (Oxadiazil 80 % 0.10 kg ha<sup>-1</sup> or Butachlor 50 EC 1.5 kg ha<sup>-1</sup>)
- 7) Hand weeding

•Seed & varieties / hybrids

- 1) Use foundation / certified seed of recommended varieties Viz., 1) KJT-2 2) KJT-3, 3) KJT-5 4) KJT-7 5) RTN-5 6) KJT-8 7) KJT-9 8) RTN-1 9) RTN-24 10) Sahyadri 11) Sahyadri-4
- 2) Seed treatment with Thirum 2.5 g kg<sup>-1</sup>
- 3) Increase area under hybrid rice upto 25 to 30 % area.
- 4) Double cropping : Cultivation of pulses, Vegetable during rabi/ summer season.

•Preparation of field: Ploughing, harrowing, puddling

- 1) Application of FYM 7.5 t ha<sup>-1</sup>
- 2) Incorporation of Dhaincha green manuring (*Sesbania rostrata* variety TSR-1)

•Transplanting: Transplanting at 20 x 15 cm or 15 x 15 cm spacing on leveled field

- 1) Age of seedling 20-25 days (1/5<sup>th</sup> of crop duration )
- 2) Application Urea DAP briquettes
- 3) Incorporation of Glyricidia leaves 5 t ha<sup>-1</sup>

•Fertilizer Management:

- 1) Integrated Nutrient management (INM)
- 2) 50 kg N, 50 kg P<sub>2</sub>O<sub>5</sub>, 50 kg K<sub>2</sub>O ha<sup>-1</sup> at transplanting
- 3) Top dressing Nitrogen 25 N at tillering & 25 kg N at panicle initiation stage

•Weed management:

- 1) Integrated Weed Management (IWM)
- 2) Spray pre-emergence weedicide Butachlor 50 EC 1.5 kg ha<sup>-1</sup>
- 3) Use of conoweeder 2 times (15 & 30 DAT)
- 4) Bunds cleaned

•Water Management :

- 1) Integrated Water Management
- 2) Maintain water level 2.5 cm for 7 days from transplanting
- 3) 5 cm upto tillering and milk stage of crop
- 4) 10 cm from panicle initiation to grain maturity
- 5) Intermittent drainage of excess water.
- 6) Drain the water 10 days before harvesting

•Pest & disease Management: Integrated pest management (IPM)

•Recommended package of practices be followed for control of

- 1) Stem borer
- 2) Brown plant hopper
- 3) Blast
- 4) BLB

•Harvesting, Threshing, Storage :

- 1) Timely mechanical harvesting.
- 2) Mechanization
- 3) Rice Processing for value addition
- 4) Minimizing the losses during pre and post harvest.
- 5) Creation of storage facilities at village level.
- 6) Extension and expansion of technological information through different extension education programmes.
- 7) Increase in minimum support price.
- 8) Adopation of Integrated farming system approach.

### Technology adoption and yield gap

The primary data on technology adoption and other economic aspects of rice production were collected from 240 randomly selected farmers from Konkan region.

The technology adoption index was estimated as

$$TAI = \frac{1}{k} \left[ \frac{AX_1}{RX_1} + \frac{AX_2}{RX_2} + \dots + \frac{AX_k}{RX_k} \right] \times 100$$

Where,

TAI = Technology adoption index

k = number of technologies

AX<sub>i</sub> = Actual use of selected technology

RX<sub>i</sub> = Recommended level of selected technology

Based on technology adopted by individual farmer sample farmers were categorised as,

## Doubling Farmers' Income (DFI) through DBSKKV Interventions

Low adopters = Mean – SD

Medium adopter = Mean – SD to mean + SD

High adopters = Mean + SD

The yield gap for different adopters category was estimated by following methodology given by IRRI, Manila, Philippines as -

$$\text{Yield gap I} = Y_p - Y_d$$

$$\text{Yield gap II} = Y_d - Y_a$$

$$\text{Total yield gap} = Y_p - Y_a$$

Where,

$Y_p$  = Potential yield realized at research station

$Y_d$  = Yield realized at demonstration plots

$Y_a$  = Yield realized on sample farms

### Distribution of sample farmers

The selected farmers were categorized into three groups as presented in Table 8. The distribution of farmers according to technology adoption index revealed that about 22 per cent farmers were high adopters, 64 per cent medium and 14 per cent low adopters. This implied that at overall level there was medium adoption of technology by the sample farmers.

### Technology adoption

Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli has released ten production technologies of rice. The technologywise level of adoption by sample farmers is given in Table 9. Table 9 revealed that, average adoption index among low, medium and high adopters was 47.91, 56.00 and 66.78 per cent, with overall average

**Table 8.** Distribution of farmers according to technology adoption.

| Sr. No. | Caterory of technology adoption | Range of Adoption Index | No. of farmers (N = 240) | %     |
|---------|---------------------------------|-------------------------|--------------------------|-------|
| 1       | Low                             | 0 to 48                 | 33                       | 13.75 |
| 2       | Medium                          | 49 to 62                | 155                      | 64.58 |
| 3       | High                            | Above 63                | 52                       | 21.67 |
| 4       | Overall adoption index          |                         | 57.22                    |       |

of 57.22 per cent. Among the different technologies of rice production adoption of harvesting technology i.e. harvesting at 90% maturity and use of Vaibhav sickle was the highest (90.67%) followed by tillage operations i.e. ploughing after harvest of previous crop at *Wafsa* condition (83.35%) and seed technology i.e. use of recommended varieties (78.81%). The adoption of technologies like use of FYM/green manuring and Urea briquette was low.

### Yield gap

The results of yield gap analysis are presented in Table 10. The yield gap analysis showed that there is inverse relation between adoption index and yield gap. It is evident that there is huge potential to increase the yield through technology adoption. The rice farmers need to be motivated to use modern rice production technologies which would add to their farm income.

### Economics of rice cultivation

The profitability of rice production among different categories of adopters is depicted in Table 11.

It is evident from Table 5 that, total cost as well as gross income increases with increase in level of adoption. The increase in cost of cultivation on high adopters farm over low adopters was 21.68 per cent. However, there was

**Table 9.** Technologywise level of adoption

| Sr. No. | Name of Technology        | Low (N=33) | Medium (N=155) | High (N=52) | Overall (N=240) |
|---------|---------------------------|------------|----------------|-------------|-----------------|
| 1.      | Tillage operations        | 62         | 85             | 92          | 83.35           |
| 2.      | F.Y.M./Green Manure       | 18         | 24             | 58          | 30.54           |
| 3.      | Seed Technology           | 61.78      | 79.54          | 87.45       | 78.81           |
| 4.      | Weedicide (for nursery)   | 0          | 0              | 0           | 0.00            |
| 5.      | Transplanting             | 66.8       | 71.33          | 84.32       | 73.52           |
| 6.      | Fertilizers               | 69.25      | 74.35          | 86.46       | 76.27           |
| 7.      | Urea Briquette            | 2.54       | 6.42           | 16.82       | 8.14            |
| 8.      | Intercultural Operations  | 46.32      | 54.48          | 69.46       | 56.60           |
| 9.      | Water level               | 67.64      | 74.64          | 77.52       | 74.30           |
| 10.     | Harvesting                | 84.72      | 90.24          | 95.72       | 90.67           |
|         | Average level of adoption | 47.91      | 56.00          | 66.78       | 57.22           |

## Special Section

reduction in per unit cost of production in high adoptor group over low adoptors by 4.94 per cent. The increase in the cost could be attributed to increase in input cost in case of high adoptor group. The benefit cost ratio revealed that rice production is marginally profitable for high adoptors. The adoption index of high adoptor group was 66.78 per cent. This implies that still there is potential to increase adoption which would ultimately results in further reduction in per unit cost of production and thereby increase in profitability in rice production.

### Prices

Price play a vital role in making agriculture remunerative. In recent past, prices of agricultural produce have become more volatile. Prices of agricultural produce are influenced by political factors, international prices, and economic factors. An attempt was made to assess the growth in farm harvest price, minimum support price and actual price realised by farms for the period from 2000-01 to 2011-12.

The Table 12 revealed that at current and constant price, farm harvest price grew at the rate of 9.69 and 3.91 per cent per annum, respectively. During the same period, growth in MSP and actual price realised by farmers at current price was 7.84 and 4.64 per cent per annum. During the same period, growth in MSP was non-significant, whereas, actual price realised by farmers, though non significant showed dealing trend. The foregoing analysis implied that net price realised by farmers for rice did not commensurate with inflation in the economy, which could be the root cause of distress in agriculture in general and rice producers in particular.

### Conclusion

To realise the goal of doubling income of farmers in Konkan region, it is necessary to identify sources of growth in income. Increase in the productivity *via* high level of technology adoption would be the significant contributor to output growth and thereby income. This requires emphasis on motivating farmers to adopt modern technologies like seed, fertilizers etc. There is considerable yield gap in rice. This gap could be bridged through technological interventions. The actual price realised by farmers for rice do not commensurate with inflation in the economy, which needs to be addressed through effective policies, and market information.

### Success Story

Shri. Vinay Maruti Vekhande a progressive farmer from Vadap village in Karjat taluka of Raigad district. Shri. Vekhande was growing traditional local rice and there after varieties like Ratna, Jaya from that he was getting yield 40 to 45 q ha<sup>-1</sup>. by adopting varieties released by Univeristy his yield was upto 50-70 q ha<sup>-1</sup>. and by

**Table 10.** Yield gap in rice (Figures in parentheses are per cent yield gap to research station yield)

| Sr. No. | Particulars                  | Low   |                  | High  | Overall          |
|---------|------------------------------|-------|------------------|-------|------------------|
| A.      | Research station yield       |       | 50.00            |       |                  |
| B.      | Demonstration plot yield     |       | 40.00            |       |                  |
| C.      | Actual yield on sample farms | 27.92 | 31.56            | 35.74 | 31.97            |
|         | Yield gap – I (A-B)          | 10.00 | 10.00            | 10.00 | 10.00            |
|         | Yield gap – II (B-C)         | 12.08 | 8.44             | 4.26  | 8.03             |
|         | Total yield gap – I (A-C)    | 22.08 | 18.44<br>(36.88) | 14.26 | 18.03<br>(36.06) |

**Table 11.** Economics of rice production

| Sr. No. | Particulars                    | Low      | Medium   | High     | Overall  |
|---------|--------------------------------|----------|----------|----------|----------|
| 1.      | Yield q ha <sup>-1</sup>       | 27.92    | 31.56    | 34.74    | 31.97    |
| 2.      | Gross income (₹)               | 40903.71 | 46390.76 | 52898.15 | 47046.22 |
| 3.      | Cost C (₹)                     | 42349.35 | 46916.02 | 51533.83 | 47288.63 |
| 4.      | Per quintal cost of production | 1516.81  | 1486.56  | 1441.91  | 1479.16  |
| 5.      | Net profit                     | -144.64  | -525.26  | 1364.32  | -242.16  |
| 6.      | B-C ratio                      | 0.97     | 0.99     | 1.03     | 0.99     |

**Table 12.** Growth in price

| Sr. No. | Particulars                     | Current price | Constant price (Base year 2004-05) |
|---------|---------------------------------|---------------|------------------------------------|
| 1.      | Farm harvest price              | 9.69**        | 3.91**                             |
| 2.      | Minimum support price           | 7.84*         | 2.16 <sup>NS</sup>                 |
| 3.      | Actual price realised by farmer | 4.64*         | -0.88 <sup>NS</sup>                |

## Doubling Farmers' Income (DFI) through DBSKKV Interventions

growing Sahyadri hybrids yield reached upto 135 q ha<sup>-1</sup>. which doubled his income.

He has adopted different techniques developed by Dr. Balasaheb Sawant Konkan Krishi Vidhyapeeth and near about doubled his income. He has followed the following techniques.

- 1) Used high yielding hybrids developed by University i.e Sahyadri hybrids and received awards for high rice yield.
- 2) Used Urea briquettes to ensure nutrient supply to the crop for long time during its growth stage.
- 3) Seed Production of different crop varieties viz Karjat-3, Karjat-7, Sahyadri hybrids, Lablab bean (Konkan wal-2) released by DBSKKV from 2012 to 2017 under Gram Bijotpadan
- 4) Used reaper for harvesting of rice and adopted mechniazation.

Shri Vekhande received appreciation from Dr. Sudhir Goel, Agriculture Commissioner, Maharashtra state and Dr. Vijay Mehta, Vice Chancellor, DBSKKV, Dapoli for seed production efforts. In 2010, he received Raigad district progressive farmer award.

During rabi season, Shri. Vekhande cultivates watermelon (hybrid) on half hectare area and Lablab

bean (Konkan wal-2) on one hectare area from which he earns good monetary returns.

### Way forward

The population is increasing day by day and land is shrinking. Therefore, rice production has to increase for maintaining food security. For rice production major constraint is labour problem at critical stages of crop growth. Therefore, mechanization will solve the problem. Reorienting research and technology development will meet emerging challenges. The future priorities are development of high value nutritionally rich varieties, tolerant to submergence, low light intensity, salt tolerant and varieties suitable for processing industry. The technology for doubling farmers income will be developed.

### References

- Anonymous 2009. Rice in India, Source : <http://Valavanstores.com>
- Anonymous 2016. All India Annual Rice Workshop Progress Report held during April 2016 at Jorhat.
- Anonymous 2017. World Agricultural Production USDA, AS, Circular Series WAP 06-17, June 2017.
- Anonymous 2017a. Maharashtra State Annual Rice Workshop Progress Report held during 10th march 2017 at pune..
- Birthal P. S., Joshi P. K., Negi D. S. and Agarwal S. 2014. Changing sources of Growth in Indian Agriculture : Implications for Regional Priorities for Accelerating Agricultural Growth. IFPRI Discussion Paper 1325, IFPRI, Washington D.C.
- Chand R., Kumar P. and Kumar S. 2011. Total Factor Productivity and Contribution of Research Investment to Agricultural Growth in India. Policy Paper 25. National Centre for Agricultural Economics and Policy Research, New Delhi.
- Dodke S. B., Shinde A. K., Torne S. R. and Mandavkar P. M. 2016. Research Recommendations 2001-2015. Directorate of Research, DBSKKV, Dapoli.
- Evenson R. E., Pray C. and Rosegrant M. W. 1999. Agricultural

**Table 13.** Production, productivity and income of Shri Vekhande prior and after adoption of DBSKKV interventions for rice.

| Year    | Yield<br>(q ha <sup>-1</sup> ) | Awards                           |
|---------|--------------------------------|----------------------------------|
| 2003-04 | 120                            | First prize in Karjat Taluka     |
| 2004-05 | 130                            | First prize in Raigad District   |
| 2005-06 | 135                            | First prize in Maharashtra State |

**Table 14.** Production, productivity and income of Shri Vekhande prior and after adoption of DBSKKV interventions for rice.

| Varieties                             | Prior  |        |        | Present  |          |                               |
|---------------------------------------|--------|--------|--------|----------|----------|-------------------------------|
|                                       | Local  | Ratna  | Jaya   | Karjat-3 | Karjat-7 | Sahyadri                      |
| Production (q)                        | 93     | 120    | 135    | 150      | 156      | 240                           |
| Productivity<br>(q ha <sup>-1</sup> ) | 31     | 40     | 45     | 50       | 52       | Potential: 120<br>Average: 80 |
| Income ha <sup>-1</sup>               | 46,500 | 60,000 | 67,500 | 75,000   | 78,000   | 1,20,000                      |

## Special Section

---

- Research and Productivity Growth in India. Research Report No. 109. IFRI, Washington D.C.
- Government of India 2017. Report of the Committee on Doubling Farmers' Income, Volume II, Status of Farmers' Income : Strategies for Accelerated Growth. Ministry of Agriculture and Farmers' Welfare, New Delhi.
- Government of India (Various issues) Commission on Agricultural Costs and Prices, New Delhi.
- Government of Maharashtra (2015-16) Socio Economic Survey of Maharashtra, Directorate of Economics and Statistics, Planning Department, Mumbai.
- Jadhav B. B., Dodke S. B., Deorukhkar A. C. and Shinde A. K. 2010. Research Accomplishments and Recommendations 2004-2009. Directorate of Research, Dr. B.S.K.K.V, Dapoli.
- Jambhale N. D., Thaware B. L. and Apte U. B. 2004. Crop varieties developed by Dr. BSKKV, Dapoli. Directorate of Research, Dr. B.S.K.K.V., Dapoli.
- Kunkerkar R. L., Vanve P. B., Waghmode B. D., Apte V. B and Gawai M. P. 2014a. High yielding medium slender rice culture KJT-2-244-10 Eco. Env & cons. 20: 349-352.
- Kunkerkar R. L., Vanve P. B., Waghmode B. D., Thaware B. L and Bhav S. G. 2014b. Super fine rice variety karjat-8 for low land mean of maharashtra state. Eco. Env. L Cons. 20: 353-356.
- Nene Y. L. 2005. Rice research on South Asia through ages Asian Agri-History 9: 85-106.
- Prasad Rao V. 2004. Role of Directorate of Rice Research in genetic improvement of rice varieties of India. Pages 101-140 in Sharma, S. D. And Prasad Rao V.(eds), Genetic Improvement of Rice varieties of India Today and Tomorrows Printers and publishers, New Delhi.
- Saxena R., Singh N. P., Balaji S. J., Ahuja U., Kumar R. and Joshi D. 2017. Doubling Farmers' Income in India by 2022-23: Sources of Growth and Approaches Agricultural Economics Research Review. Vol. 30(2), pp. 265-277.
- Sehgal J. L. Mandal D. K. Mandal C and Vedinelu S. 1992. Agroecological Regions of India . Technical Bulletin No.24 2nd Edition NBSS & LUp, ICAR Nagpur, Maharashtra, India, 130.
- Sharma G. R. 1985. from hunting and food gathering to domestication of plants and animals in the Belan and Ganga Valleys Pages 369-371 in Misra V. N and Bellwood, P. (eds), Recent Advantages in Indo-Pacific Prehistory, Oxford and IBH, New Delhi.
- Shoba Rani, N. Prasad, G. S. V. 2008. High Yielding rice Varieties of India. Technical Bulletin No.33. Directorate of Rice Research, Hyderabad.
- Swarna S. V., Mofarahus S. and Sharma S. D. 2010. History of Rice in South Asia (1947-2007). In: Sharma S. D. (ed). Rice: Origin Antiquity and History, Taylor and Francis. pp 272-283.
-